

Medication

Your dog has been provided with a complimentary oral suspension for pain relief. This is to be given for a few days after surgery with food.

If she has any vomiting or diarrhoea please stop this medication and contact your veterinary surgery.

Post-operative checks

You will need to make an appointment for post-operative checks 3 days and 10 days after surgery, or as advised on discharge.

If you are worried in between appointments, please contact The Pet Vet at any time.



The Pet Vet Doncaster
25-29 Balby Road, Doncaster, DN4 ORD
Tel: 01302 325608
Email: doncaster@thepetvet.co.uk

The Pet Vet Rotherham 158 Doncaster Road, Dalton, Rotherham, S65 3EW Tel: 01709 854333

Email: rotherham@thepetvet.co.uk

The Pet Vet Barnsley Wakefield Road, Barnsley, S71 1NN

Tel: 01226 297125 Email: barnsley@thepetvet.co.uk


The Pet Vet Lincoln
Unit 8A, Carlton Boulevard, Carlton Centre,
Lincoln. LN2 4FJ

Tel: 01522 458895 Email: lincoln@thepetvet.co.uk



2022-08-DOG

thepetvet.co.uk





Your dog has been spayed today, which involves the surgical removal of her ovaries and womb in sterile conditions. Now that she is home, here are a few things you can do to make her feel more comfortable.

She has experienced a general anaesthetic, and so she may appear quiet or sleepy for the next 24 hours; this is to be expected, and she should be back to normal after this time.



We recommend gastrointestinal food (available from the surgery) to be given for 48 hours post-operatively, which helps the digestive system after the anaesthetic. Please make sure she has access to fresh water at all times.

If she doesn't like this diet, you can offer her something light such as boiled white fish, chicken or boiled rice. Don't worry if your pet doesn't respond to food straightaway; they don't always want to eat after an anaesthetic.



Your dog needs restricted exercise for up to 10 days after their operation.

Dogs should be limited to short lead exercise only until you are advised otherwise and you should restrict her from jumping up and down as much as possible.

Stitches

It is very important that your dog cannot lick or chew her wound.

We advise you to use, where possible, a buster collar or medical pet shirt to prevent interference.